

Drug	Strength of tab (mg) or liquid mg/ml	Number of tablets or ml by weight band (twice daily)													Strength of adult tab (mg)	Number of tablets by weight band (twice daily)	
		Children 6 weeks of age and above (0.75 BD is delivered as 1 tablet AM and 0.5 tablets PM and 1.5 BD is delivered as 2 tablets AM and 1 tablet PM)														25-29.9 kg	30-34.9 kg
		3-3.9 kg	4-4.9 kg	5-5.9 kg	6-6.9 kg	7-7.9 kg	8-8.9 kg	9-9.9 kg	10-10.9 kg	11-11.9 kg	12-13.9 kg	14-16.9 kg	17-19.9 kg	20-24.9 kg			
AZT	60	1	1	1	1.5	1.5	1.5	1.5	2	2	2	2.5	2.5	3	300	1	1
AZT (new annex E)	300; 10 mg/ml	6 ml	6 ml	6 ml	9 ml	9 ml	9 ml	9 ml	12 ml	12 ml	12 ml	0.5	0.5	0.75	300	1	1
AZT/3TC	60/30	1	1	1	1.5	1.5	1.5	1.5	2	2	2	2.5	2.5	3	300/150	1	1
AZT/3TC/NVP	60/30/50	1	1	1	1.5	1.5	1.5	1.5	2	2	2	2.5	2.5	3	300/150/200	1	1
ABC	60	1	1	1	1.5	1.5	1.5	1.5	2	2	2	2.5	2.5	3	300	1	1
ABC (new annex E)	300; 20 mg/ml	3 ml	3 ml	3 ml	4 ml	4 ml	4 ml	4 ml	6 ml	6 ml	6 ml	0.5	0.5	0.75	300	1	1
ABC/3TC	60/30	1	1	1	1.5	1.5	1.5	1.5	2	2	2	2.5	2.5	3	300/150	1	1
ABC/3TC/NVP	60/30/50	1	1	1	1.5	1.5	1.5	1.5	2	2	2	2.5	2.5	3	300/150/200	1	1
ABC/AZT/3TC	60/60/30	1	1	1	1.5	1.5	1.5	1.5	2	2	2	2.5	2.5	3	300/300/150	1	1
3TC	30	1	1	1	1.5	1.5	1.5	1.5	2	2	2	2.5	2.5	3	150	1	1
3TC (new annex E)	150; 10 mg/ml	3 ml	3 ml	3 ml	4 ml	4 ml	4 ml	4 ml	6 ml	6 ml	6 ml	0.5	0.5	0.75	150	1	1
d4T	6	1	1	1	1.5	1.5	1.5	1.5	2	2	2	2.5	2.5	3	30	1	1
d4T (new annex E)	various; 1 mg/ml	6 ml	6 ml	6 ml	9 ml	9 ml	9 ml	9 ml	1x15 mg	1x15 mg	1x15 mg	1x20 mg	1x20 mg	1x20 mg	30	1	1
d4T/3TC	6/30	1	1	1	1.5	1.5	1.5	1.5	2	2	2	2.5	2.5	3	30/150	1	1
d4T/3TC/NVP	6/30/50	1	1	1	1.5	1.5	1.5	1.5	2	2	2	2.5	2.5	3	30/150/200	1	1
NVP	50	1	1	1	1.5	1.5	1.5	1.5	2	2	2	2.5	2.5	3	200	1	1
NVP (new annex E)	200; 10 mg/ml	5 ml	5 ml	5 ml	8 ml	8 ml	8 ml	8 ml	10 ml	10ml	10 ml	0.75	0.75	0.75	200	1	1
Lopinavir/ritonavir	100/25	n/r	n/r	n/r	n/r	n/r	n/r	n/r	1.5	1.5	1.5	2	2	2.5	100/25* (paed)	3	3
Lop/rit (new annex E)	80/20 mg/ml	1 ml	1.5 ml	1.5 ml	1.5 ml	1.5 ml	1.5 ml	1.5 ml	2 ml	2 ml	2 ml	2.5 ml	2.5 ml	3 ml	80/20 mg/ml	3.5 ml	4 ml

* 3 tablets BD of 100/25 may be substituted with 2 tablets am and 1 tablet pm of 200/50

Note: higher doses of Lop/rit may be required when co-administered with enzyme-inducing drugs such as NVP, EFV; fosamprenavir, rifampicin.